



Technical data sheet Serum

Ref : FT.SxxxFan
Page : 1/3

Version date : 23/08/2022

Fetal Bovine Serum Charcoal Stripped

CAT N°: SxxxF

Collected from the source:

When searchers choose their serum an important factor that should be taken into consideration is the source, which also emphasises the traceability of the serum.

Our system of vertical integration allows us to be certain of the origins and traceability of our FBS.

Each manufactured batch is rigorously controlled, from the collection of serum and throughout all stages of its treatment and production through to final packaging on our premises.

BioWest Fetal Bovine Serum is derived from clotted whole blood aseptically collected from foetus via cardiac puncture.

The serum is collected or imported and treated in agreement with the European regulations.

Country of Origin:

The country in which the serum was taken from the donor/animal.

To see the countries of origin we can offer, please refer to the technical data sheet for the standard Fetal Bovine Serum (ref. FT.FBSan).

To order the treated serum, please replace the last number of the Cat N° of standard serum (written on the FT.FBSan) by the letter F.

Storage conditions:

- 18°C to - 40°C, protected from light.

Bottles can be stored between -40°C and -80°C for a short period (few days).

Shelf life:

5 years

Filtration:

Final Filter Size: 0.2µm

pH:

pH specification: Not applicable

Osmolality:

Determined by a lowered freezing temperature. The osmometer is calibrated against standard solutions.

Osmolality specification: Not applicable

Endotoxin:

All sera are tested to determine the levels of endotoxins. BioWest carries out a chromokinetic quantitative test, according to the method D of the European Pharmacopoeia.

The endotoxin reagent is standardized against the US reference endotoxin.

Endotoxin specification: As low as possible

Haemoglobin:

The haemoglobin level is measured by spectrophotometer.

Haemoglobin specification: As low as possible

Cell Culture:

Biological performance is assessed using cell culture medium supplemented with the serum being tested.

During the test period, cultures are examined microscopically for any morphological abnormalities that may indicate toxic components in the serum.



Technical data sheet Serum

Ref : FT.SxxxFan
Page : 2/3

Version date : 23/08/2022

Cell Culture Tests:

Cell Growth, Plating Efficiency, Cloning Efficiency.

Cell Lines Tested:

The following cell lines are tested with the serum:

- HELA -Cancer Cell/Human.
- L929 -Fibroblast-Mouse/ As Macrophage
- SP2/O-AG14 -Mouse/Lymphoma
- MRC- 5 -Human/Lung.

Total Protein:

Determined by Biuret Colorimetry.

Total protein specification: Not applicable

Sterility tests:

All sera are tested for the absence of aerobic and anaerobic bacteria, fungi, yeast and *Mycoplasma*.

The sterility test is based on the European Pharmacopoeia requirements.

The sera are tested for the absence of *Mycoplasma* by culture.

Virus test:

All of our sera are tested for:

- Bovine Viral Diarrhoea (BVD)
- Cytopathogenic agents e.g. Infectious Bovine Rhinotracheitis (IBR) / BHV-1
- Hemadsorbing agents e.g. Parainfluenza Type 3 (PI3)

Sera are tested by inoculation to permissive cells. The revelation is made by immunofluorescence for pestiviruses. Cytopathogenic agents and hemadsorbing agents are detected by microscopic observations.

Other tests:

The concentration of following hormones is measured before and after treatment by radioimmunology:

- Oestradiol
- Progesteron
- Testosteron

Hormones specification: Not applicable

Treatments:

Charcoal stripped serum is treated by filtering chilled serum through an activated carbon adsorbent filter to remove non-polar material. This treatment removes lipophilic material but has little effect on the concentration of salts dissolved in the serum.

BioWest offers Charcoal / Dextran stripped foetal bovine serum for researchers requiring low levels of various hormones. Charcoal / Dextran stripping reduces the concentration of steroid hormones in serum eg estradiol, progesterone, cortisol, testosterone, T3 and T4. It has been demonstrated and used in numerous studies both *in vivo* and *in vitro* (e.g., 1,2,3,4). So this serum is useful for utilisation in receptor studies, oestrogen related investigations, or when endogenous steroid hormones may interfere with experimental work.

In addition, Charcoal / Dextran treatment has been demonstrated to improve immunoassay systems (5,6,7,8,9,10,11); Herbert, et al. (12) showed that the use of Charcoal / Dextran improved insulin assay methods.

Moreover, some studies indicate that Charcoal / Dextran treatments may be used to minimise lot to lot serum variability.

This serum may show some reduced growth promotion of cells requiring the presence of certain hormones and growth factors.

The level of endotoxin is in general higher than non treated serum because of the endotoxins in the charcoal and in the dextran.



Technical data sheet Serum

Ref : FT.SxxxFan
Page : 3/3

Version date : 23/08/2022

Recommended use:

- Respect storage conditions of the serum
- Do not use the serum after its expiry date
- Store serum in an area protected from light
- Manipulate serum in aseptic conditions (e.g.: under laminar air flow)
- Wear clothes adapted to the manipulation of serum to avoid contamination (e.g.: gloves, mask, hygiene cap, overall...)
- In order to preserve all serum qualities, it is recommended to thaw out the flask, to aliquote, then to re-freeze the produced flasks rather than to thaw out and re-freeze the flask at each use.
- It is recommended to use the serum immediately after its thaw out. However, if it is not useful, it is possible to store thaw out serum, at +2°C / +8°C, until 26 weeks without significant decrease of its performances in cell culture.

The product is intended to be used *in vitro* for research or further manufacturing only and not for use as an Active Pharmaceutical Ingredient or food or animal feed.

Remarks:

The raw serum may be treated (Heat Inactivated, Gamma Irradiated, pH modified) before filtration for different reasons:

- Importation regulation
- Exportation necessity
- Technical or quality aspects.

To be informed if your batch is concerned by treatment before filtration, please contact Biowest.

REFERENCES :

1. Carson, R.S., D.M. Robertson and J.K. Findlay. 1988. Ovine follicular fluid inhibits thymidine incorporation by 3T3 fibroblasts *in vitro*. J. Reprod. Fertil. 82: 447-455.
2. Coezy, E., J. Bouhnik, E. Clauser, F. Pinet, M. Philippe, J. Menard and P. Corvol. 1984. Effects of glucocorticoids and antiglucocorticoid on angiotensinogen production by hepatoma cells in culture. *In Vitro* 20: 528-534.
3. Friedman, E.A., M.J. Saltzman, B.G. Delano and M.M. Beyer. 1978. Reduction in hyperlipidemia in hemodialysis patients treated with charcoal and oxidized starch (oxystarch). *Am. J. Clin. Nutr.* 31: 1903-1914.
4. Seaver, S.S., J. van der Bosch and G. Sato. 1984. The chick oviduct in tissue culture. I. Initial characterization of growing primary oviduct tissue cultures. *Exp. Cell Res.* 155: 241-251.
5. Croze, F. and P. Francimont. 1984. Biological determination of inhibin in rat ovarian-cell culture medium. *J. Reprod. Fertil.* 72: 237-248.
6. Ding, Y.Q. and I. Huhtaniemi. 1989. Human serum LH inhibitor(s): behaviour and contribution to *in vitro* bioassay of LH using dispersed mouse Leydig cells. *Acta Endocrinol. (Copenh.)* 121: 46-54.
7. Helin, H.J., J.J. Isola, M.J. Helle and H. Adlercreutz. 1988. Influence of endocrine status on biochemical and immunocytochemical estrogen and progesterone receptor assays in breast cancer patients. *Breast Cancer Res. Treat.* 12: 67-73.
8. Kohno, T. and E. Ishikawa. 1986. A highly sensitive enzyme immunoassay of anti-insulin antibodies in guinea pig serum. *J. Biochem. (Tokyo)* 100: 1247-1251.
9. Nakamura, S., S. Sakata, T. Komaki, K. Kamikubo, K. Yasuda and K. Miura. 1986. An improved and simplified method for the detection of thyroid hormone autoantibodies (THAA) in serum. *Endocrinol. Jpn.* 33: 415-422.
10. Silbert, C.K. and C.T. Sawin. 1975. Double-antibody radioimmunoassay of serum insulin: effect of use of hormone-depleted human serum. *Clin. Chem.* 21: 1520-1522.
11. Yoshimura, M., Y. Ochi, T. Hachiya and T. Miyazaki. 1975. Estimation of the maximal T4-binding capacity of TBG using the Triosorb test in serum treated with dextran-coated charcoal. *Endocrinol. Jpn.* 22: 199-205.
12. Herbert V., K.-S. Lau, C.W. Gottlieb and S.J. Bleicher. 1965. Coated charcoal immunoassay of insulin. *J. Clin. Endocrinology* 25:1375-1384.